## CONSENT FOR TWO-STAGE OSSEOUSINTEGRATED IMPLANT WITH SINUS-LIFT/BONE GRAFTING PROCEDURE Page 1 of 4

Patient's Na	me Date			
Please initial each paragraph after reading. If you have any questions, please ask your doctor before initialing.				
are able to to sign is	the right to be given information about your proposed implant placement so that you make the decision as to whether to proceed with surgery. What you are being asked confirmation that you have been given information on the nature of your proposed the known risks associated with it and the possible alternative treatments.			
1.	Drhas informed me of my diagnosis (condition) which is described as:			
2.	The surgical procedure proposed to treat the above condition has been explained to me and I understand it to be:			
3.	I understand that incisions will be placed inside my mouth in the upper jaw for the purpose of placing one or more endosteal root form structures (implants) in my jaw to serve as anchors for a missing tooth or teeth replacement or to stabilize a crown (cap), bridge or denture. I acknowledge that the doctor has explained the procedure, including the number and location of incisions and the type of implant to be used. I also understand that the crown, bridge or denture that will later be attached to this implant(s) will be made and attached by Dr and that a separate charge will be made by that office.			
4.	In my case, I further understand that there is not enough natural jawbone in which to place the proposed implant and that a procedure called "sinus lift" is planned. This procedure is more complicated than usual implant placement and involves opening the sinus cavity in my upper jaw and placing a bone graft in order to provide support for the implant. I have been told that this graft could come from specially-prepared donated bone, or may be taken from my jaw, chin, skull or hip, any of which might be supplemented with specially-prepared donated bone or bone substitute.			
5.	I understand the implant(s) must remain covered by gum tissue for at least six months or longer before it can be used, and that a second surgical procedure is required to uncover the top of the implant(s). No guarantee can be or has been given that the implant(s) will last for a specific time period. It has been explained to me that once the implant(s) is/are inserted, the entire treatment plan must be followed and completed on schedule. If this is not done, the implant(s) may fail.			

## CONSENT FOR TWO-STAGE OSSEOUSINTEGRATED IMPLANT WITH SINUS-LIFT/BONE GRAFTING PROCEDURE Page 2 of 4

I have be	en informed of possible alternative forms of treatment (if any), including:
	and that other forms of treatment or no treatment at all are choices that I have sks of those choices have been presented to me.
side effec	or has explained to me that there are certain inherent and potential risks and ets in any surgical procedure and in this specific instance such risks include, of limited to, the following:
RISKS C	OF IMPLANT SURGERY
A.	
B.	at-home recuperation.  Prolonged or heavy bleeding that may require additional treatment.  Because the sinus is involved, some bleeding may be from the nose.
C.	Injury or damage to adjacent teeth or roots of adjacent teeth, possibly requiring further root canal therapy, and occasionally the loss of an injured tooth.
D.	Post-operative infection, including sinus infection, which may require additional treatment. In rare instances an opening may develop between mouth and sinus, again requiring additional treatment.
E.	Stretching of the corners of the mouth that may cause cracking and bruising.
F.	Restricted mouth opening for several days; sometimes related to swelling and muscle soreness, and sometimes related to stress on the jaw joints (TMJ).
G.	Possible prolonged symptoms of sinusitis requiring certain medications and longer recovery time, resulting from intentional entry into the sinus.
H. I.	Fracture of the jaw.  Possible injury to nerve branches in the bone resulting in numbness, pair or tingling of the lips, cheek, gums or teeth. If implants are placed in the lower jaw, there may be numbness or pain of the chin or tongue also These symptoms may persist for several weeks, months or, in rare instances, may be permanent.
GENER	AL RISKS OF BONE GRAFTING
A.	Bleeding, swelling or infection at the donor site requiring further
B.	treatment.  Allergic or other adverse reaction to drugs used during or after the procedure.
C.	• • · · · · · · · · · · · · · · · · · ·

## CONSENT FOR TWO-STAGE OSSEOUSINTEGRATED IMPLANT WITH SINUS LIFT/BONE GRAFTING PROCEDURE Page 3 of 4

RISKS AND COMPLICATIONS OF GRAFTING FROM WITHIN THE

	MOUTH AREA
	<ul> <li>A. Damage to adjacent teeth, which may require future root canal procedures, or may cause loss of those teeth.</li> <li>B. Removal of adult teeth in order to obtain sufficient bone material.</li> <li>C. Numbness or pain in the area of the donor or recipient site, or more extensive areas, which may be temporary or permanent.</li> <li>D. Penetration of the sinus or nasal cavity in the upper jaw which could result in infection or other complication requiring additional drug or surgical treatment.</li> </ul>
	RISKS OF FREEZE-DRIED, DEMINERALIZED OR OTHER BANKED BONE
	On occasion, additional donated bone is used to supplement the patient's bone, or to spare an extensive donor site surgical procedure. Use of such bone may involve separate risks including, but not limited to:
	A. Rejection of the donated graft material together with the entire graft.  B. The remote chance of disease transmission from processed bone.
8.	I understand that in my grafting procedure, the use of (autogenous, demineralized, etc.) bone is expected to be taken from (note anatomic area), plus(other area)
9.	ANESTHESIA The anesthesia I have chosen for my surgery is:     Local Anesthesia     Local Anesthesia with Nitrous Oxide/Oxygen Analgesia     Local Anesthesia with Oral Premedication     Local Anesthesia with Intravenous Sedation     General Anesthesia
10.	ANESTHETIC RISKS include: discomfort, swelling, bruising, infection, prolonged numbness and allergic reactions. There may be inflammation at the site of an intravenous injection (phlebitis) which may cause prolonged discomfort and/or disability, and may require special care. Nausea and vomiting, although rare, may be unfortunate side effects of IV anesthesia. Intravenous anesthesia is a serious medical procedure and, although considered safe, carries with it the risk of heart irregularities, heart attack, stroke, brain damage or death.

## CONSENT FOR TWO-STAGE OSSEOUSINTEGRATED IMPLANT WITH SINUS LIFT/BONE GRAFTING PROCEDURE Page 4 of 4

11.	YU	OUR OBLIGATIONS IF IV ANESTHESIA IS USED			
		Because anesthetic medications cause prolonged drowsiness, you MUST be accompanied by a responsible adult to drive you home and stay with you until			
		you are recovered sufficiently to care for yourself. This may be up to 24 hours.			
	B.	During recovery time (24 hours) you should not drive, operate complicated			
		machinery or devices, or make important decisions such as signing documents etc.			
	C.	You must have a completely empty stomach. <u>IT IS VITAL THAT YOU HAVE</u>			
		NOTHING TO EAT OR DRINK FOR EIGHT (8) HOURS PRIOR TO YOUR			
	D	ANESTHETIC. TO DO OTHERWISE MAY BE LIFE-THREATENING!  However, it is important that you take any regular medications (high blood			
	<b>D</b> .	pressure, antibiotics, etc.) or any medications provided by this office, using only			
		a small sip of water.			
12	)	It has been explained to me that in the course of the procedure unforeseen			
		y be revealed which will necessitate extension of the original procedure, a different			
	•	n those set forth above, or abandonment of the procedure entirely. In such an			
		ize my doctor and his staff to perform such procedures as are necessary and			
desirable i	n the	e exercise of professional judgment to complete my surgery.			
13.	I uı	nderstand that my doctor is not a seller of the implant device itself and makes no			
	wai	rranty or guarantee regarding success or failure of the implant or its attachments			
	use	d in the procedure.			
14.	I u	nderstand smoking is extremely detrimental to the success of implant surgery.			
		ee to cease all use of tobacco for 2-3 weeks prior to and after surgery, including			
		later uncovering procedure, and to make strong efforts to give up smoking			
	ent	irely.			
15.	It h	as been explained to me and I understand that a perfect result is not, and cannot be			
	guaranteed or warranted.				
CONSEN	т				
		speak, read and write English and have read and fully understand this consent for			
-		at all blanks were filled in prior to my initialing and signing this form and that all			
ny questio	ons v	were answered to my satisfaction.			
Patient's (or	Lega	al Guardian's) Signature Date			
ationt 5 (or	Logo	a Guardian of Signature			
Doctor's Sig	matui	re Date			
20001 3 018	,	Duc			
Witness' Sig	natui	re Date			